

Investigation of Oxidative Stress Defenses of *Neisseria gonorrhoeae* by Using a Human Polymorphonuclear Leukocyte Survival Assay

Kate L. Seib,¹ Mark P. Simons,² Hsing-Ju Wu,³ Alastair G. McEwan,¹ William M. Nauseef,² Michael A. Apicella,² and Michael P. Jennings^{1*}

The School of Molecular and Microbial Sciences and The Centre for Metals in Biology, The University of Queensland, Brisbane, Australia¹; Department of Microbiology, Inflammation Program, and Department of Medicine, University of Iowa, Iowa City, Iowa²; and Core Facilities for Proteomics Research, Institute of Biological Chemistry, Academia Sinica, Taipei, Taiwan³

Received 24 January 2005/Returned for modification 4 March 2005/Accepted 21 March 2005

***Neisseria gonorrhoeae* has well-characterized oxidative stress defense systems that protect against oxidative killing in in vitro assays. In contrast, mutant strains of *N. gonorrhoeae* lacking oxidative stress defenses are identical to the wild type when tested in an ex vivo survival assay using human polymorphonuclear leukocytes.**

Neisseria gonorrhoeae, the causative agent of the sexually transmitted infection gonorrhea, is a host-adapted pathogen that poses serious health implications. Gonococcal infection is typically characterized by a symptomatic inflammatory response of the urogenital tract, which involves accumulation of activated polymorphonuclear leukocytes (PMNs) (4). The PMN-mediated inflammatory response involves migration of PMNs towards sites of infection, phagocytosis of microorganisms, and elimination of these organisms by oxygen-dependent and oxygen-independent mechanisms (reviewed in references 11 and 23).

Stimulation of PMNs results in a rapid increase in oxygen consumption referred to as the “oxidative burst.” Assembly and activation of the NADPH oxidase in the plasma membrane results in the generation of superoxide (5), which rapidly dismutates to hydrogen peroxide (27) and is then consumed by myeloperoxidase, generating hypochlorous acid (52). Nitric oxide is also produced by constitutive and inducible nitric oxide synthases of PMNs (12, 25, 28). Various secondary oxidants are generated from these reactive species, including chloramines, hydroxyl radicals, singlet oxygen, and peroxynitrite (12, 23). Concurrent with the oxidative burst, intracellular granules fuse with the plasma membrane or phagosomal membrane to release a broad array of biologically active molecules, including proteases and antimicrobial proteins (reviewed in reference 18).

Without antibiotic treatment, gonococcal infections are persistent and resolve slowly (33), indicating that the PMN response is relatively ineffective in eradicating infection. However, interaction of *N. gonorrhoeae* with PMNs has been a controversial topic; some studies have reported that *N. gonorrhoeae* cells survive and grow within PMNs, while other studies have reported that *N. gonorrhoeae* is rapidly killed within PMNs (reviewed in references 37 and 44). The majority of these studies were performed using a tumbling tube assay with nonadherent PMNs. The use of adherent PMNs mimics the natural disease process in which PMNs have migrated from the

bloodstream and are attached to cells and extracellular matrix proteins present at the site of infection. PMNs adhered to plates coated with serum or extracellular matrix proteins produce a large respiratory burst in response to stimuli, whereas suspension PMNs are relatively unresponsive (32). A recent study investigating interactions between *N. gonorrhoeae* and adherent human PMNs found the PMNs generated a substantial respiratory burst in response to gonococci (45). Despite this, a significant proportion of phagocytosed *N. gonorrhoeae* cells survived PMN killing and replicated over time (45). Viable counts and microscopic analysis indicated that some level of killing occurs after ingestion, but a subpopulation of *N. gonorrhoeae* cells survive and replicate (75.0% ± 18.31% at 1 h and 80.21% ± 15.34% at 2 h), in contrast to efficient killing of *Escherichia coli* (1.83% ± 0.36% at 1 h and 1.37% ± 0.08% at 2 h) (45).

It seemed probable that the ability of *N. gonorrhoeae* cells to survive in the hostile environment of the PMN would be due to the diverse array of oxidative stress defenses that this organism possesses. These defenses have typically been characterized based on the sensitivity of mutant strains to in vitro oxidative killing (see Table 1 for a summary). *N. gonorrhoeae* possesses one superoxide dismutase (Sod), which is an inactive or weakly active cytoplasmic SodB (4, 24, 53) that does not protect against reactive oxygen species in vitro (49). However, accumulation of manganese by the MntABC transport system confers protection against oxidative stress (49). *N. gonorrhoeae* also has high catalase and peroxidase activities (4, 24, 53), afforded by the cytoplasmic catalase (KatA) (4) and the periplasmic cytochrome *c* peroxidase (Ccp), both of which protect against hydrogen peroxide in vitro (42, 50). Other oxidative defenses described for *N. gonorrhoeae* include the following: a potential thiol-disulfide oxidoreductase, Sco (41); methionine sulfoxide reductase (MsrA/B) (46); and the iron storage protein bacterioferritin (Bfr) (14). Regulators of these defenses include the peroxide-responsive repressors of KatA and MntC, OxyR (48) and PerR (H.-J. Wu, K. L. Seib, Y. N. Srikhanta, J. L. Edwards, S. P. Kidd, M. A. Apicella, A. G. McEwan, and M. P. Jennings, submitted for publication), respectively.

* Corresponding author. Mailing address: The School of Molecular and Microbial Sciences, The University of Queensland, Brisbane 4072, Australia. Phone: 61 7 3365 4879. Fax: 61 7 3365 4620. E-mail: jennings@uq.edu.au.

TABLE 1. Strains used in the PMN phagocytosis assay and a summary of results

Strain	Description of strain	In vitro sensitivity ^a		PMN sensitivity ^c
		Sensitivity (substance)	Reference	
<i>N. gonorrhoeae</i> 1291	Wild type			—
<i>sodB</i>	Sod deficient	X (PQ, X/XO)	49	—
<i>mntC</i>	Lacks periplasmic Mn-binding protein of MntABC transporter (decreased [Mn] in cells)	S (PQ)	49	—
<i>katA</i>	Catalase deficient	S (H ₂ O ₂)	42	—
<i>ccp</i>	Cytochrome <i>c</i> peroxidase deficient	S (H ₂ O ₂ [slight])	42	—
<i>ccp katA</i>	Ccp and catalase double mutant	S (H ₂ O ₂)	42	—
<i>sco</i>	Sco deficient (potential thiol-disulfide oxidoreductase)	S (PQ)	41	—
<i>sco katA</i>	Sco and catalase double mutant	S (PQ, H ₂ O ₂)	41	—
<i>oxyR</i>	Overexpresses KatA	R (X/XO, H ₂ O ₂)	48	—
<i>perR</i>	Overexpresses MntC (increased [Mn] in cells)	R (H ₂ O ₂)	— ^b	—
<i>E. coli</i> DH5α	Wild type			+

^a The strains used in the PMN phagocytosis assay are shown along with a summary of the results from this assay and results from in vitro assays from previous publications. Sensitivities of mutant strains to in vitro oxidative stress killing assays are shown relative to *N. gonorrhoeae* strain 1291 (wild type): X, same phenotype as wild type; S, sensitive to killing; R, resistant to killing (increased survival relative to wild type). These assays involved exposure of a suspension of 10⁴ to 10⁶ cells to either paraquat (PQ; 10 mM), xanthine (4.3 mM)/xanthine oxidase (300 mU/ml) (X/XO), or hydrogen peroxide (H₂O₂; 10 or 40 mM). For further details, see the references cited in the table.

^b —, H.-J. Wu, K. L. Seib, Y. N. Srihanta, J. L. Edwards, S. P. Kidd, M. A. Apicella, A. G. McEwan, and M. P. Jennings, submitted for publication.

^c PMN sensitivity (+) is defined as a statistically significant difference in the mean percent survival of the strain relative to *N. gonorrhoeae* strain 1291 (wild type) in the PMN phagocytosis assay. *P* values that are <0.05 are considered statistically significant. Differences between *N. gonorrhoeae* 1291 and its mutant derivatives were not statistically significant (—) (Fig. 1) (*P* values ≥ 0.14).

The oxidative stress that *N. gonorrhoeae* encounters within the host environment is significantly more complex than that generated by in vitro assays. Indeed, the expression of numerous genes involved in defense against oxidative stress is up-regulated in *E. coli* upon exposure to PMNs (47). Many of these defenses (i.e., catalase, Sod, and OxyR) have been shown to protect enteric bacteria against killing by activated PMNs (19, 29, 39, 47). The role of oxidative defenses in protecting *N. gonorrhoeae* from the PMN oxidative burst was determined using the phagocytosis assay described by Simons et al. (45). Briefly, approximately 2 × 10⁶ PMNs isolated from peripheral blood were adhered to glass coverslips pretreated with collagen and autologous human serum. *N. gonorrhoeae* strain 1291 (wild type) and a set of mutant derivatives (Table 1) expressing opacity-associated protein (Opa⁺) and pili (P⁺) (as determined by colony morphology) were grown to log phase. Bacteria were added to PMNs at a multiplicity of infection of 1:1 and centrifuged, and the supernatant was removed (time A). Synchronized phagocytosis was stimulated by the addition of warm medium for 2 minutes (time B). Nonphagocytosed bacteria were removed by three wash steps (time 0 h). PMNs and bacteria were incubated at 37°C for 1 hour (time 1 h). Saponin was used to lyse PMNs, and viable CFU were enumerated after overnight culture of plated serial dilutions. Each assay was done with triplicate cultures of each mutant strain along with wild-type controls and was performed on at least three occasions. Laser scanning confocal microscopy, scanning electron microscopy, and transmission electron microscopy performed by Simons et al. (45) illustrated that PMN-associated *N. gonorrhoeae* cells were internalized and contained within defined phagosomes within the perimeter of the PMN membrane and that there were very few bacteria residing in the extracellular spaces.

This study is based on a set of previously constructed and defined mutant strains of *N. gonorrhoeae* that are deficient in various oxidative stress defense mechanisms or regulatory systems (see Table 1 and references cited therein for further

detail), with the exception of the *sco katA* mutant strain, which was constructed by transforming the *sco::kan* plasmid (41) into the *katA::tet* strain (42). All mutant strains were confirmed by Southern hybridization, PCR, and/or sequence analysis. The set of mutant strains was compared to wild-type strain 1291 to determine relative sensitivity to killing by PMNs (Fig. 1). A summary of results is shown in Table 1, along with results of in vitro assays from previously published work. Surprisingly, none of the mutant strains tested had a phenotype distinct from the wild type in the PMN phagocytosis assay. Wild-type and mutant strains had similar levels of PMN association (time A) and phagocytosis (time B), indicating the mutations had no detectable effect on phagocytosis by PMNs.

N. gonorrhoeae is a host-adapted pathogen which is able to survive killing by PMNs. Based on the broad range of oxidative stress defenses that *N. gonorrhoeae* cells possess, it was proposed that the ability to survive within PMNs was dependent, at least in part, on protection against the oxidative burst. This hypothesis is not supported by the similar phenotypes that were observed for the wild-type strain and the oxidative stress defense mutant strains in the PMN assay.

Several species of bacteria are able to survive PMN killing by inhibiting or subverting the oxidative burst of PMNs (reviewed in reference 1). As seen with *N. gonorrhoeae* in this study, *Helicobacter pylori* is able to evade PMN killing despite the activation of an oxidative response. *H. pylori* alters the targeting of the PMN NADPH oxidase so that it locates to the plasma membrane rather than the phagosomal membrane, thus releasing reactive oxygen species into the extracellular environment (2).

The features that govern *N. gonorrhoeae*-PMN interactions have not yet been defined, and while interactions do involve the gonococcal surface proteins porin (8, 21, 22, 26, 31) and Opa (7, 20, 31, 51), their roles in stimulating versus inhibiting the oxidative burst of PMNs have not been resolved. Further investigation using the adherent PMN phagocytosis assay (45) should provide a better understanding of the *N. gonorrhoeae*-




FIG. 1. PMN phagocytosis assay. PMNs were incubated with *N. gonorrhoeae* wild-type strain 1291 and mutant derivatives deficient in various oxidative stress defense mechanisms or regulatory systems to determine relative sensitivity to killing by PMNs (as described by Simons et al. [45]). Briefly, bacteria were added to a monolayer of PMNs (approximately 2×10^6) at a multiplicity of infection of 1:1 and incubated at 37°C for 1 h. PMN were then lysed to quantify CFU. Each panel (A to G) shows a set of experiments in which wild-type strain 1291 (Ng) was compared to a mutant strain, a set of mutant strains, or *Escherichia coli* strain DH5 α (Ec), as shown on the x axis. Results shown are the percent survival of bacteria at time 1 h (expressed as a percentage of viable bacteria at time 0 h). Each data point is the mean of several individual experiments \pm one standard deviation of the mean (number of repeat experiments: panel A, 3; B, 4; C, 2; D, 4; E, 2; F, 3; and G, 3) in which each experiment was conducted with three replicates of each strain. To determine whether the percent survival of mutant strains was statistically significant compared to the percent survival of the wild-type strain, *P* values were computed using unpaired two-sided Student's *t* test. *P* values that were less than 0.05 were considered statistically significant. Differences between *N. gonorrhoeae* wild-type strain 1291 and its mutant derivatives were not statistically significant (panels A to F, *P* \geq 0.14). The only statistically significant difference was between wild-type *N. gonorrhoeae* and *E. coli* (panel G; *P* < 0.0001).

dependent oxidative burst of PMNs. Despite the stimulation of an oxidative burst by *N. gonorrhoeae* (7, 31, 45, 51), the results described herein support previous studies which suggested that oxygen-independent mechanisms may be of greater significance than oxygen-dependent mechanisms during PMN killing of *N. gonorrhoeae* (9, 10, 13, 35, 36, 43; reviewed in reference 44).

The resistance of the specific mutant strains may also be due to the presence of redundant defenses. For example, in *E. coli*, a *sodA* or a *sodB* mutant strain is no more sensitive to PMN killing than the wild-type strain, but a *sodA sodB* double mutant is more susceptible to killing (29, 34). It is also important to note that the defenses investigated do not protect against all of the types of oxidants generated during the PMN oxidative burst (e.g., nitric oxide), some of which may be of greater importance in the killing of *N. gonorrhoeae*.

In light of our results, the presence of a wide range of defenses in *N. gonorrhoeae* suggests that this organism encounters significant oxidative stress from other sources in vivo. The primary sites of gonococcal infection are the ecto- and endocervical epithelium in women (17) and the urethral epithelium in men (3, 15). *N. gonorrhoeae* cells are able to survive and replicate within epithelial cells at these sites of infection (reviewed in reference 30). Intestinal and airway epithelial cells are able to kill bacteria by oxidative mechanisms (6, 16, 38, 40), and cervical epithelial cells also appear to have such oxidative defense capacity. We have recently observed that both MntC and PerR of *N. gonorrhoeae* are required for survival within primary human cervical epithelial cells (H.-J. Wu, K. L. Seib, Y. N. Srikhanta, J. L. Edwards, S. P. Kidd, M. A. Apicella, A. G. McEwan, and M. P. Jennings, submitted for publication).

This work was supported by Program Grant 284214 from the National Health and Medical Research Council of Australia and by U.S. Public Health service grants AI45728, AI43924, and AI38515 from the National Institute of Allergy and Infectious Diseases. K.L.S. acknowledges The University of Queensland Graduate School for providing a Graduate School Research Travel Award.

REFERENCES

- Allen, L. A. 2003. Mechanisms of pathogenesis: evasion of killing by polymorphonuclear leukocytes. *Microbes Infect.* 5:1329–1335.
- Allen, L. A. 2000. Modulating phagocyte activation: the pros and cons of *Helicobacter pylori* virulence factors. *J. Exp. Med.* 191:1451–1454.
- Apicella, M. A., M. Ketterer, F. K. Lee, D. Zhou, P. A. Rice, and M. S. Blake. 1996. The pathogenesis of gonococcal urethritis in men: confocal and immunoelectron microscopic analysis of urethral exudates from men infected with *Neisseria gonorrhoeae*. *J. Infect. Dis.* 173:636–646.
- Archibald, F. S., and M. N. Duong. 1986. Superoxide dismutase and oxygen toxicity defenses in the genus *Neisseria*. *Infect. Immun.* 51:631–641.
- Babior, B. M., J. D. Lambeth, and W. Nauseef. 2002. The neutrophil NADPH oxidase. *Arch. Biochem. Biophys.* 397:342–344.
- Battistoni, A., F. Pacello, S. Folcarelli, M. Ajello, G. Donnarumma, R. Greco, M. G. Ammendolia, D. Touati, G. Rotilio, and P. Valenti. 2000. Increased expression of periplasmic Cu,Zn superoxide dismutase enhances survival of *Escherichia coli* invasive strains within nonphagocytic cells. *Infect. Immun.* 68:30–37.
- Belland, R. J., T. Chen, J. Swanson, and S. H. Fischer. 1992. Human neutrophil response to recombinant neisserial Opa proteins. *Mol. Microbiol.* 6:1729–1737.
- Bjerknes, R., H. K. Guttormsen, C. O. Solberg, and L. M. Wetzler. 1995. Neisserial porins inhibit human neutrophil actin polymerization, degranulation, opsonin receptor expression, and phagocytosis but prime the neutrophils to increase their oxidative burst. *Infect. Immun.* 63:160–167.
- Britigan, B. E., and M. S. Cohen. 1986. Effects of human serum on bacterial competition with neutrophils for molecular oxygen. *Infect. Immun.* 52:657–663.
- Britigan, B. E., D. Klapper, T. Svendsen, and M. S. Cohen. 1988. Phagocyte-derived lactate stimulates oxygen consumption by *Neisseria gonorrhoeae*. *An*

- unrecognized aspect of the oxygen metabolism of phagocytosis. *J. Clin. Investig.* **81**:318–324.
11. Burg, N. D., and M. H. Pillinger. 2001. The neutrophil: function and regulation in innate and humoral immunity. *Clin. Immunol.* **99**:7–17.
 12. Carreras, M. C., G. A. Pargament, S. D. Catz, J. J. Poderoso, and A. Boveris. 1994. Kinetics of nitric oxide and hydrogen peroxide production and formation of peroxynitrite during the respiratory burst of human neutrophils. *FEBS Lett.* **341**:65–68.
 13. Casey, S. G., W. M. Shafer, and J. K. Spitznagel. 1986. *Neisseria gonorrhoeae* survive intraleukocytic oxygen-independent antimicrobial capacities of anaerobic and aerobic granulocytes in the presence of pyocin lethal for extracellular gonococci. *Infect. Immun.* **52**:384–389.
 14. Chen, C. Y., and S. A. Morse. 1999. *Neisseria gonorrhoeae* bacterioferritin: structural heterogeneity, involvement in iron storage and protection against oxidative stress. *Microbiology* **145**:2967–2975.
 15. Cohen, M. S., J. G. Cannon, A. E. Jerse, L. M. Charniga, S. F. Isbey, and L. G. Whicker. 1994. Human experimentation with *Neisseria gonorrhoeae*: rationale, methods, and implications for the biology of infection and vaccine development. *J. Infect. Dis.* **169**:532–537.
 16. Deitch, E. A., Y. Haskel, N. Cruz, D. Xu, and P. R. Kvietys. 1995. Caco-2 and IEC-18 intestinal epithelial cells exert bactericidal activity through an oxidant-dependent pathway. *Shock* **4**:345–350.
 17. Edwards, J. L., J. Q. Shao, K. A. Ault, and M. A. Apicella. 2000. *Neisseria gonorrhoeae* elicits membrane ruffling and cytoskeletal rearrangements upon infection of primary human endocervical and ectocervical cells. *Infect. Immun.* **68**:5354–5363.
 18. Faurschou, M., and N. Borregaard. 2003. Neutrophil granules and secretory vesicles in inflammation. *Microbes Infect.* **5**:1317–1327.
 19. Franzon, V. L., J. Arondel, and P. J. Sansonetti. 1990. Contribution of superoxide dismutase and catalase activities to *Shigella flexneri* pathogenesis. *Infect. Immun.* **58**:529–535.
 20. Gray-Owen, S. D., D. R. Lorenzen, A. Haude, T. F. Meyer, and C. Dehio. 1997. Differential Opa specificities for CD66 receptors influence tissue interactions and cellular response to *Neisseria gonorrhoeae*. *Mol. Microbiol.* **26**:971–980.
 21. Haines, K. A., J. Reibman, X. Y. Tang, M. Blake, and G. Weissmann. 1991. Effects of protein I of *Neisseria gonorrhoeae* on neutrophil activation: generation of diacylglycerol from phosphatidylcholine via a specific phospholipase C is associated with exocytosis. *J. Cell Biol.* **114**:433–442.
 22. Haines, K. A., L. Yeh, M. S. Blake, P. Cristello, H. Korchak, and G. Weissmann. 1988. Protein I, a translocatable ion channel from *Neisseria gonorrhoeae*, selectively inhibits exocytosis from human neutrophils without inhibiting O_2^- generation. *J. Biol. Chem.* **263**:945–951.
 23. Hampton, M. B., A. J. Kettle, and C. C. Winterbourn. 1998. Inside the neutrophil phagosome: oxidants, myeloperoxidase, and bacterial killing. *Blood* **92**:3007–3017.
 24. Johnson, S. R., B. M. Steiner, and G. H. Perkins. 1996. Cloning and characterization of the catalase gene of *Neisseria gonorrhoeae*: use of the gonococcus as a host organism for recombinant DNA. *Infect. Immun.* **64**:2627–2634.
 25. Kudoh, S., K. Suzuki, M. Yamada, Q. Liu, S. Nakaji, and K. Sugawara. 1999. Contribution of nitric oxide synthase to human neutrophil chemiluminescence. *Luminescence* **14**:335–339.
 26. Lorenzen, D. R., D. Gunther, J. Pandit, T. Rudel, E. Brandt, and T. F. Meyer. 2000. *Neisseria gonorrhoeae* porin modifies the oxidative burst of human professional phagocytes. *Infect. Immun.* **68**:6215–6222.
 27. Makino, R., T. Tanaka, T. Iizuka, Y. Ishimura, and S. Kanegasaki. 1986. Stoichiometric conversion of oxygen to superoxide anion during the respiratory burst in neutrophils. Direct evidence by a new method for measurement of superoxide anion with diacetyldeuterioheme-substituted horseradish peroxidase. *J. Biol. Chem.* **261**:11444–11447.
 28. McCall, T. B., N. K. Boughton-Smith, R. M. Palmer, B. J. Whittle, and S. Moncada. 1989. Synthesis of nitric oxide from L-arginine by neutrophils. Release and interaction with superoxide anion. *Biochem. J.* **261**:293–296.
 29. McManus, D. C., and P. D. Josephy. 1995. Superoxide dismutase protects *Escherichia coli* against killing by human serum. *Arch. Biochem. Biophys.* **317**:57–61.
 30. Merz, A. J., and M. So. 2000. Interactions of pathogenic neisseriae with epithelial cell membranes. *Annu. Rev. Cell Dev. Biol.* **16**:423–457.
 31. Naidu, F. L., and R. F. Rest. 1991. Stimulation of human neutrophil oxidative metabolism by nonopsonized *Neisseria gonorrhoeae*. *Infect. Immun.* **59**:4383–4390.
 32. Nathan, C. F. 1987. Neutrophil activation on biological surfaces. Massive secretion of hydrogen peroxide in response to products of macrophages and lymphocytes. *J. Clin. Investig.* **80**:1550–1560.
 33. Ober, W. B. 1970. Boswell's clap. *JAMA* **212**:91–95.
 34. Papp-Szabo, E., C. L. Sutherland, and P. D. Josephy. 1993. Superoxide dismutase and the resistance of *Escherichia coli* to phagocytic killing by human neutrophils. *Infect. Immun.* **61**:1442–1446.
 35. Rest, R. F. 1979. Killing of *Neisseria gonorrhoeae* by human polymorphonuclear neutrophil granule extracts. *Infect. Immun.* **25**:574–579.
 36. Rest, R. F., S. H. Fischer, Z. Z. Ingham, and J. F. Jones. 1982. Interactions of *Neisseria gonorrhoeae* with human neutrophils: effects of serum and gonococcal opacity on phagocyte killing and chemiluminescence. *Infect. Immun.* **36**:737–744.
 37. Rest, R. F., and W. M. Shafer. 1989. Interactions of *Neisseria gonorrhoeae* with human neutrophils. *Clin. Microbiol. Rev.* **2**(Suppl.):S83–S91.
 38. Rochelle, L. G., B. M. Fischer, and K. B. Adler. 1998. Concurrent production of reactive oxygen and nitrogen species by airway epithelial cells in vitro. *Free Radic. Biol. Med.* **24**:863–868.
 39. Rosen, H., P. J. Lewis, and C. M. Nitzel. 2004. Neutrophil microbicidal activity: screening bacterial mutants for survival after phagocytosis using quantitative PCR. *Jpn. J. Infect. Dis.* **57**:S19–S21.
 40. Schmidt, H. H., and U. Walter. 1994. NO at work. *Cell* **78**:919–925.
 41. Seib, K. L., M. P. Jennings, and A. G. McEwan. 2003. A Sco homologue plays a role in defence against oxidative stress in pathogenic *Neisseria*. *FEBS Lett.* **546**:411–415.
 42. Seib, K. L., H. J. Tseng, A. G. McEwan, M. A. Apicella, and M. P. Jennings. 2004. Defenses against oxidative stress in *Neisseria gonorrhoeae* and *Neisseria meningitidis*: distinctive systems for different lifestyles. *J. Infect. Dis.* **190**:136–147.
 43. Shafer, W. M., V. C. Onunka, and L. E. Martin. 1986. Antagonococcal activity of human neutrophil cathepsin G. *Infect. Immun.* **54**:184–188.
 44. Shafer, W. M., and R. F. Rest. 1989. Interactions of gonococci with phagocytic cells. *Annu. Rev. Microbiol.* **43**:121–145.
 45. Simons, M. P., W. M. Nauseef, and M. A. Apicella. 2005. Interactions of *Neisseria gonorrhoeae* with adherent polymorphonuclear leukocytes. *Infect. Immun.* **73**:1971–1977.
 46. Skaar, E. P., D. M. Tobiason, J. Quick, R. C. Judd, H. Weissbach, F. Etienne, N. Brot, and H. S. Seifert. 2002. The outer membrane localization of the *Neisseria gonorrhoeae* MsrA/B is involved in survival against reactive oxygen species. *Proc. Natl. Acad. Sci. USA* **99**:10108–10113.
 47. Staudinger, B. J., M. A. Oberdoerster, P. J. Lewis, and H. Rosen. 2002. mRNA expression profiles for *Escherichia coli* ingested by normal and phagocyte oxidase-deficient human neutrophils. *J. Clin. Investig.* **110**:1151–1163.
 48. Tseng, H. J., A. G. McEwan, M. A. Apicella, and M. P. Jennings. 2003. OxyR acts as a repressor of catalase expression in *Neisseria gonorrhoeae*. *Infect. Immun.* **71**:550–556.
 49. Tseng, H. J., Y. Srihanta, A. G. McEwan, and M. P. Jennings. 2001. Accumulation of manganese in *Neisseria gonorrhoeae* correlates with resistance to oxidative killing by superoxide anion and is independent of superoxide dismutase activity. *Mol. Microbiol.* **40**:1175–1186.
 50. Turner, S. M., E. G. Reid, H. Smith, and J. A. Cole. 2003. A novel cytochrome c peroxidase from *Neisseria gonorrhoeae*, a lipoprotein from a Gram-negative bacterium. *Biochem. J.* **373**:865–873.
 51. Virji, M., and J. E. Heckels. 1986. The effect of protein II and pili on the interaction of *Neisseria gonorrhoeae* with human polymorphonuclear leukocytes. *J. Gen. Microbiol.* **132**:503–512.
 52. Winterbourn, C. C., M. C. Vissers, and A. J. Kettle. 2000. Myeloperoxidase. *Curr. Opin. Hematol.* **7**:53–58.
 53. Zheng, H. Y., D. J. Hassett, K. Bean, and M. S. Cohen. 1992. Regulation of catalase in *Neisseria gonorrhoeae*. Effects of oxidant stress and exposure to human neutrophils. *J. Clin. Investig.* **90**:1000–1006.